

# Composite Likelihood Methods for Large Bayesian VARs with Stochastic Volatility

Joshua Chan<sup>1</sup>   Eric Eisenstat<sup>2</sup>   Chenghan Hou<sup>3</sup>   Gary  
Koop<sup>4</sup>

<sup>1</sup>University of Technology Sydney   <sup>2</sup>University of Queensland   <sup>3</sup>Hunan University  
<sup>4</sup>University of Strathclyde

# Background: History of Large VARs

- Large VARs, involving 100 or more dependent variables, are increasingly used in a variety of macroeconomic applications.
- Pioneering paper: Banbura, Giannone and Reichlin (2010, JAE) "Large Bayesian Vector Autoregressions"
- Previous VARs: a few variables perhaps 10 at most
- BGR has 131 variables (standard US macro variables)
- Many others, here is a sample:
- Carriero, Kapetanios and Marcellino (2009, IJF): exchange rates for many countries
- Carriero, Kapetanios and Marcellino (2012, JBF): US government bond yields of different maturities
- Giannone, Lenza, Momferatou and Onorante (2010): euro area inflation forecasting (components of inflation)
- Koop and Korobilis (2016, EER) eurozone sovereign debt crisis
- Bloor and Matheson (2010, EE): macro application for New Zealand
- Jarociński and Maćkowiak (2016, ReStat): Granger causality

# Background: Why large VARs?

- Availability of more data
- More data means more information, makes sense to include it
- Concerns about missing out important information (omitted variables bias, fundamentalness, etc.)
- The main alternatives are factor models
- Principal components squeeze information in large number of variables to small number of factors
- But this squeezing is done without reference to explanatory power (i.e. squeeze first then put in regression model or VAR): “unsupervised”
- Large VAR methods are supervised and can easily see role of individual variables
- And they work: often beating factor methods in forecasting competitions

# Background: Computation in large VARs

- E.g. large VAR with  $N = 100$  variables and a lag length of  $p = 13$ :
- 100,000+ VAR coefficients
- 5,050 free parameters in error covariance.
- Bayesian prior shrinkage surmounts over-parameterization
- Standard choices exist: e.g. Minnesota prior
- Key point 1: Standard approaches are conjugate: analytical results exist (estimation and forecasting – no MCMC needed)
- Key point 2: Huge posterior covariance of VAR coefficients ( $N^2 p \times N^2 p$  matrix): tough computation
- Key point 3: Conjugacy greatly simplifies: separately manipulate  $N \times N$  and  $Np \times Np$  matrices
- Key point 4: Using more realistic priors or extending model (e.g. to relax homoskedasticity assumption) loses conjugacy and, thus, computational feasibility
- Bottom line: Great tools exist for large homoskedastic Bayesian VARs with a particular prior, but cannot easily

# Background: Multivariate Stochastic Volatility in VARs

- Allowing for error variances to change in macroeconomic VARs important
- E.g. Primiceri (2005, ReStud), Sims and Zha (2006, AER), Clark (2011, JBES), etc.
- Research question: How to add multivariate stochastic volatility in large VARs?
- Existing Bayesian literature is either:
  - Homoskedastic
  - Restrictive forms (e.g. Clark, Carriero and Marcellino, 2016, JBES + 2 working papers, Chan, 2016, working paper)
  - Approximations (Koop and Korobilis, JOE, JOE and Koop, Korobilis and Pettenuzzo, 2016, JOE)
- Present paper: new approach using composite likelihoods

# Vector Autoregressions with Stochastic Volatility (VAR-SV)

- $y_t$  is  $N$ -vector of dependent variables ( $N$  large)
- VAR-SV is:

$$A_{0t}y_t = c + A_1y_{t-1} + \dots + A_p y_{t-p} + \epsilon_t, \quad \epsilon_t \sim N(0, \Sigma_t),$$

- $\Sigma_t = \text{diag}(e^{h_{1,t}}, \dots, e^{h_{n,t}})$

- 

$$A_{0t} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ a_{21,t} & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1,t} & a_{n2,t} & \dots & 1 \end{pmatrix}$$

- Rewrite as

$$y_t = X_t\beta + W_t a_t + \epsilon_t$$

- $X_t = I_n \otimes (1, y'_{t-1}, \dots, y'_{t-p})$
- $a_t$  is vector of free elements of  $A_{0t}$

# Vector Autoregressions with Stochastic Volatility

- $W_t = \begin{pmatrix} 0 & 0 & 0 & \cdots & \cdots & \cdots & 0 \\ -y_{1,t} & 0 & 0 & \cdots & \cdots & \cdots & 0 \\ 0 & -y_{1,t} & -y_{2,t} & \cdots & \cdots & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots & \cdots & \cdots & 0 \\ 0 & \cdots & \cdots & -y_{1,t} & -y_{2,t} & \cdots & -y_{N-1,t} \end{pmatrix}$

- 

$$h_t = h_{t-1} + \epsilon_t^h, \quad \epsilon_t^h \sim N(0, \Sigma_h)$$

$$a_t = a_{t-1} + \epsilon_t^a, \quad \epsilon_t^a \sim N(0, \Sigma_a)$$

- $\Sigma_h = \text{diag}(\sigma_{h,1}^2, \dots, \sigma_{h,N}^2)$  and  $\Sigma_a = \text{diag}(\sigma_{a,1}^2, \dots, \sigma_{a, \frac{N(N-1)}{2}}^2)$ .
- Standard MCMC methods used for estimation and forecasting
- But these will not work with large VARs

# Composite Bayesian Methods

- Likelihood function (assuming independent errors):

$$L(y; \theta) = \prod_{t=1}^T p(y_t | \theta) = \prod_{t=1}^T L(y_t; \theta)$$

- Composite likelihood

$$L^C(y; \theta) = \prod_{t=1}^T \prod_{i=1}^M L^C(y_{i,t}; \theta)^{w_i}$$

- $y_{i,t}$  for  $i = 1, \dots, M$  are sub-vectors of  $y_t$
- $L^C(y_{i,t}; \theta) = p(y_{i,t} | \theta)$
- $w_i$  weight attached to sub-model  $i$
- $\sum_{i=1}^M w_i = 1$
- Bayesian composite posterior

$$p^C(\theta | y) \propto L^C(y; \theta) p(\theta)$$



# How do we use composite Bayesian methods?

- Instead of forecasting with large VAR-SV, forecast with many small VAR-SVs
- Let  $y_t = \begin{pmatrix} y_t^* \\ z_t \end{pmatrix}$
- $y_t^*$  contains  $N_*$  variables of interest
- $z_t$  (with elements denoted by  $z_{i,t}$ ) remaining variables.
- Sub-model  $i$  is VAR-SV using  $y_{i,t} = \begin{pmatrix} y_t^* \\ z_{i,t} \end{pmatrix}$
- Our application uses 193 variables with  $N_* = 3$
- Thus, 190 sub-models, each is a 4-variate VAR-SV

- Some asymptotic theory exists (e.g. Canova and Matthes, 2017)
- Require strong assumptions
- Overview: Varin, Reid and Firth (2011, Stat Sin)
- Pakel, Shephard, Sheppard and Engle (2014, working paper)
- Need asymptotic mixing assumptions about dependence over time, over variables and between different variables at different points in time
- In general, strong assumptions often not achieved in practice
- Hence, our justification is mostly empirical

# Theory of Composite Likelihoods as Opinion Pools

- Bayesian theory uses idea of opinion pool
- Each sub-model is “agent” with “opinion” about a feature (e.g. a forecast) expressed through a probability distribution.
- Theory addresses “How do we combine these opinions?”
- Generalized logarithmic opinion pool equivalent to composite likelihood
- Nice properties (e.g. external Bayesianity)
- Linear opinion pools lead to other combinations of sub-models
- E.g. Geweke and Amisano (2011, JOE) optimal prediction pools
- In empirical work consider both composite likelihood and Geweke-Amisano

# Choosing the Weights

- Various approaches considered
- Equal weights  $w_i = \frac{1}{M}$
- Weights proportional to marginal likelihood of each sub-model
- Weights proportional to (exponential of) BIC of each sub-model
- Weights proportional to (exponential of) DIC of each sub-model
- In all above use likelihood/marginal likelihood for core variables only ( $y_t^*$ )

- Target: Draws from Bayesian composite posterior

$$p^C(\theta|y) \propto L^C(y; \theta) p(\theta)$$

- We have:
  - 1. MCMC draws from  $M$  sub-models (4-variate VAR-SVs)
  - 2. Weights,  $w_i$  for  $i = 1, \dots, M$
- We develop accept-reject algorithm
- See paper for details

# Macroeconomic Forecasting Using a Large Dataset

- FRED-QD data set from 1959Q1- 2015Q3
- 193 quarterly US variables (transformed to stationarity)
- Three core variables: CPI inflation, GDP growth and the Federal Funds rate.
- Small data set: 7 variables
- Core variables + unemployment, industrial production, money (M2) and stock prices (S&P)
- Large data set: All 193 variables
- Lag length of 4

- With small data set use variety of models
- Computation is feasible (and over-parameterization concerns smaller)
- Large data set:
- Compare composite likelihoods methods to homoskedastic, conjugate prior, large VAR

- For composite likelihood approach prior elicitation less of an issue (small models)
- With large VARs prior elicitation is crucial (may or may not be disadvantage)
- For all models use comparable priors
- Hyperparameter choices inspired by Minnesota prior
- See paper for details



- Variety of different weights in composite likelihood approaches
- Standard VAR-SV (Primiceri, 2005, ReStud)
- Homoskedastic VARs of different dimensions
- Carriero, Clark and Marcellino (CCM, 2016a,b)
- CCM1: common drifting volatility model
- VAR-SV with  $a_t = 0$  and  $\Sigma_t = e^{h_t} \Sigma$
- $h_t$  is scalar stochastic volatility process
- CCM2: more flexible SV model
- VAR-SV with  $a_t$  constant
- Each equation error has own volatility, but restrictions on correlations

---

---

	Description
VAR-HM	7-variable Homoskedastic VAR
VAR-SV	7-variable VAR with stochastic volatility
VAR-CCM1	7-variable model of CCM (2016a)
VAR-CCM2	7-variable model of CCM (2016b)
Large VAR	large Homoskedastic VAR
VAR-CL-BIC	VAR-CL-SV with BIC based weights
VAR-CL-DIC	VAR-CL-SV with DIC based weights
VAR-CL-EQ	VAR-CL-SV with equal weights
VAR-GA	VAR-SV with G-A weights
VAR-CL-ML	VAR-CL-SV with ML weights


---

---


# Estimating Variances and Covariances

- Key variables of interest (common to all models) are  $\sigma_{ij,t}$  for  $i, j = 1, 2, 3$
- Small data set: VAR-SV will probably be closest to “true” specification (most flexible)
- Evaluate performance relative to VAR-SV
- VAR-SV in red in following figures
- Dotted lines in some figures credible intervals (16th-84th percentiles)


# Estimating Variances and Covariances




# Comparison of VAR-CL-ML to VAR-SV




# Comparison of VAR-CCM1 to VAR-SV



# Comparison of VAR-CCM2 to VAR-SV



# Comparison of VAR-HM to VAR-SV





- Estimation results are encouraging, what about forecasting?
- Results for  $h = 1$
- Two forecast evaluation periods:
- Beginning 1970Q1
- Beginning 2008Q1 (financial crisis and subsequent recession)

- For 3 core variables individually:
- RMSFE
- MAFE
- ALPL = average of log predictive likelihoods (higher value better)
- ACRPS = average of conditional rank probability score (lower values better)
- Also joint ALPL based on joint predictive for core variables

# Joint ALPL for Core Variables

Forecast Performance	Post-1970	Post-2008
VAR-HM	0.33	-0.58
VAR-SV	0.65	0.44
VAR-CCM1	0.06	-0.51
VAR-CCM2	0.90	0.52
Large VAR	-0.47	-1.69
VAR-CL-ML	0.90	<b>1.27</b>
VAR-CL-DIC	0.85	0.67
VAR-CL-BIC	0.90	1.15
VAR-CL-EQ	0.88	0.89
VAR-GA	<b>0.91</b>	1.01

- Best overall summary
- Composite likelihoods + Geweke-Amisano forecast best
- Weights: Marginal likelihood or BIC weights best (but only slightly)
- Homoskedastic large VAR does poorly
- CCM2 better than CCM1

# Forecasting the Core Forecasts Individually

- Following tables present results for each variable
- General themes:
- Composite likelihoods+GA forecast well
- Especially for 2008-2016 period
- Especially for inflation and interest rate
- Less so for GDP growth (VAR-SV is best)
- Large homoskedastic VARs forecast poorly
- CCM2 better than CCM1
- In general, CCM2 similar but a bit worse than composite likelihoods

# Inflation Forecasting Beginning in 1970

	RMSFE	MAE	ACRPS	ALPL
VAR-HM	<b>0.66</b>	<b>0.45</b>	0.36	-0.15
VAR-SV	0.67	0.46	0.36	-0.06
VAR-CCM1	0.71	0.51	0.39	-0.12
VAR-CCM2	0.67	0.46	0.36	-0.00
Large VAR	0.73	0.52	0.56	-0.14
VAR-CL-ML	0.69	0.47	0.36	-0.01
VAR-CL-DIC	0.68	0.47	0.36	-0.01
VAR-CL-BIC	0.69	0.46	0.36	-0.01
VAR-CL-EQ	0.68	0.47	0.36	-0.01
VAR-GA	0.68	0.47	0.38	<b>-0.00</b>

# Inflation Forecasting Beginning in 2008

	RMSFE	MAE	ACRPS	ALPL
VAR-HM	1.04	0.66	0.52	-1.16
VAR-SV	1.06	0.68	0.54	-0.68
VAR-CCM1	1.04	0.66	0.52	-0.71
VAR-CCM2	1.05	0.68	0.53	-0.57
Large VAR	1.03	0.65	0.69	-0.71
VAR-CL-ML	1.04	0.65	0.51	-0.54
VAR-CL-DIC	1.04	0.66	0.52	-0.57
VAR-CL-BIC	<b>1.02</b>	<b>0.63</b>	<b>0.50</b>	-0.50
VAR-CL-EQ	1.04	0.66	0.52	-0.57
VAR-GA	1.03	0.66	0.54	<b>-0.48</b>

# Interest Rate Forecasting Beginning in 1970

	RMSFE	MAE	ACRPS	ALPL
VAR-HM	0.29	0.18	0.15	0.81
VAR-SV	0.28	0.17	0.14	1.03
VAR-CCM1	0.51	0.33	0.25	0.53
VAR-CCM2	0.28	0.17	0.14	1.19
Large VAR	0.56	0.42	0.44	0.17
VAR-CL-ML	0.28	0.17	0.13	1.18
VAR-CL-DIC	0.28	0.16	0.13	1.17
VAR-CL-BIC	0.28	0.17	0.13	1.20
VAR-CL-EQ	0.27	0.16	0.13	1.19
VAR-GA	<b>0.27</b>	<b>0.16</b>	0.13	<b>1.21</b>



# Interest Rate Forecasting Beginning in 2008

	RMSFE	MAE	ACRPS	ALPL
VAR-HM	0.25	0.18	0.14	0.97
VAR-SV	0.18	0.12	0.10	1.50
VAR-CCM1	0.36	0.30	0.20	0.66
VAR-CCM2	0.20	0.12	0.10	1.45
Large VAR	0.51	0.45	0.46	0.09
VAR-CL-ML	0.13	<b>0.07</b>	<b>0.06</b>	<b>2.00</b>
VAR-CL-DIC	0.13	0.08	0.07	1.67
VAR-CL-BIC	0.13	0.07	0.06	1.88
VAR-CL-EQ	0.12	0.08	0.07	1.79
VAR-GA	<b>0.12</b>	0.08	0.07	1.83

# GDP growth Forecasting Beginning in 1970

	RMSFE	MAE	ACRPS	ALPL
VAR-HM	0.89	0.68	0.51	-0.38
VAR-SV	<b>0.86</b>	<b>0.65</b>	<b>0.50</b>	-0.32
VAR-CCM1	0.87	0.67	0.51	-0.36
VAR-CCM2	0.86	0.66	0.50	<b>-0.31</b>
Large VAR	0.93	0.70	0.77	-0.39
VAR-CL-ML	0.92	0.67	0.51	-0.35
VAR-CL-DIC	0.91	0.67	0.51	-0.36
VAR-CL-BIC	0.93	0.68	0.52	-0.35
VAR-CL-EQ	0.92	0.68	0.51	-0.35
VAR-GA	0.92	0.68	0.54	-0.36

# GDP growth Forecasting Beginning in 2008

	RMSFE	MAE	ACRPS	ALPL
VAR-HM	0.96	0.72	0.56	-0.48
VAR-SV	<b>0.86</b>	<b>0.63</b>	<b>0.50</b>	<b>-0.42</b>
VAR-CCM1	0.94	0.73	0.57	-0.57
VAR-CCM2	0.88	0.65	0.52	-0.46
Large VAR	0.96	0.77	0.80	-0.47
VAR-CL-ML	0.95	0.65	0.52	-0.46
VAR-CL-DIC	0.95	0.66	0.53	-0.50
VAR-CL-BIC	0.96	0.67	0.52	-0.47
VAR-CL-EQ	0.95	0.66	0.52	-0.47
VAR-GA	0.96	0.68	0.56	-0.46

- Composite likelihood methods allows VAR-SV with huge data sets
- Computationally and conceptually simple: average over many small models
- Other VAR-SV models have some attractive features but are computationally infeasible with huge data sets
- In small data set, composite likelihood methods approximate other methods
- In large data set, composite likelihoods forecast better than large VAR